

Isolation, cloning and molecular analysis of *ag85a* and *tb10.4* genes from *Mycobacterium tuberculosis*

Roghayeh Teimourpour ¹, Zahra Meshkat ^{2*}, Samira Rashidian ²

¹Department of Microbiology, School of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran

²Antimicrobial Resistance Research Center, Bu Ali Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, IR Iran, E-mail: meshkatz@mums.ac.ir

Background and Objective: Novel TB vaccines that aim to boost and/or replace Bacillus Calmette-Guerin (BCG) are currently in development. DNA vaccines can stimulate both humoral and cell-mediated immunity in different animal models of TB and is thought to be a promising strategy in the development of new vaccines against TB. The aim of this study was to design and construct a DNA vaccine encoding *ag85a* and *tb10.4* fusion genes of *Mycobacterium tuberculosis*

Materials and Methods: *tb10.4* fragment was amplified by PCR and the products were digested with restriction enzymes. Next, it was cloned into the pcDNA3.1⁺ plasmid. The *ag85a* gene and pcDNA3.1⁺/*tb10.4* plasmid were digested by *EcoRI* and *BamHI* restriction enzymes. Constructed vector was sequenced and molecular analysis was done by using bioinformatic software such as DNAMAN. New chimeric vector containing *Ag85a-tb10.4* genes were purified and for confirming expression of pcDNA3.1⁺/*tb10.4-ag85a* plasmid in eukaryotic cells, Huh7.5 cell lines were transfected with this recombinant vector.

Results: Using electrophoresis of PCR products, fragments 297 bp for *tb10.4* and 1017 bp for *ag85a* were observed. alignment of *ag85a-tb10.4* genom sequence with reference genes in GenBank showed exact identity that indicate correction of all cloning procedures. Eukaryotic cells transfection with pcDNA3.1⁺/*tb10.4-ag85a* vector was confirmed with cDNA synthesis and existence of *tb10.4-ag85a* fusion gene was confirmed with RT-PCR (Fig1,2,3,4).

Conclusion:

In this study, *tb10.4* and *ag85a* genes were isolated from *Mycobacterium tuberculosis* H37Rv strain and cloned into pcDNA3.1⁺ also in this study capability of constructed vector in producing fusion *Ag85a-tb10.4* protein was confirmed with RT-PCR. pcDNA3.1⁺/*tb10.4-ag85a* vector can be used for further studies in future.




Fig.1: Purified PCR product of *tb10.4* gene on 1.5% gel : Lane 1: 290 bp band of *tb10.4*, Lane 2: 1kb DNA size marker (SM0313, Fermentas, Germany)




Figure 2: Purified PCR product of *ag85a* gene on 1.5% gel : Lane 1: 1017 bp band of *ag85a*, Lane 2: 1kb DNA size marker (SM0313, Fermentas, Germany)




Fig 3: Restriction enzyme digestion of pcDNA3.1⁺/*ag85a-tb10.4*. Lane 1: 1kb DNA size marker (SM0313, Fermentas, Germany). Lane 2: 1300 bp band of *tb10.4/ag85a*.




Fig4: RT-PCR results. Lane 1: 1300 bp band of *ag85a/tb10.4*. Lane 2: 1kb DNA size marker (SM0313, Fermentas, Germany).

References

1. Skjot RLV, Brock I, Arend SM, Munk ME, Theisen M, Ottenhoff TH, et al. Epitope mapping of the immunodominant antigen TB10. 4 and the two homologous proteins TB10. 3 and TB12. 9, which constitute a subfamily of the esat-6 gene family. *Infection and immunity*. 2002;70(10):5446-53.
2. Akhavan R, Meshkat Z, Khajekaramadini M, Meshkat M. Eight-year study of mycobacterium tuberculosis in mashhad, northeast of iran. *Iranian Journal of Pathology*. 2013;8(2):73-80.
3. Asnaashari AMH, Sadrizadeh A, Ahmadi H, Meshkat M, Gholoobi A, Talab FR, et al. The study of Mycobacterium tuberculosis in Iranian patients with lung cancer. *Jundishapur J Microbiol*. 2013;6:237-41.
4. McShane H. Tuberculosis vaccines: beyond bacille Calmette-Guérin. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2011;366(1579):2782-9.
5. Organization WH. Global tuberculosis control: WHO report 2010: World Health Organization; 2010.